



Contents lists available at ScienceDirect

Heart & Lung

journal homepage: www.heartandlung.com

Promising impacts of mesenchymal stem cell therapy in treatment of SARS-CoV-2 (COVID-19)

Masoud Khorshidi, Ph.D.^{a,b}, Meysam Zarezadeh, Ph.D.^{c,d,**}, Mohammadreza Emami, M.Sc.^e, Beheshteh Olang, MD, Ph.D.^b, Omid Moradi Moghaddam, MD, FCCM^{f,*}

^a Student Research Committee, Department of Nutrition, School of Public Health, Iran University of Medical Sciences, Tehran, Iran

^b Pediatric Gastroenterology, Hepatology and Nutrition Research Center, Research Institute for Children's Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

^c Department of Clinical Nutrition, Nutrition Research Center, School of Nutrition and Food Science, Tabriz University of Medical Sciences, Tabriz, Iran

^d Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

^e Department of Clinical Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences (TUMS), Tehran, Iran

^f Trauma and Injury Research Center, Critical Care Medicine Department, Iran University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article History:

Received 11 August 2020

Accepted 17 August 2020

Available online 19 August 2020

To the Editor

A novel coronavirus named COVID-19 has begun to spread in Wuhan, China from December 2019 and become a global health concern.¹ Individuals with underlying diseases are at greater risk and more prone to be critically ill in case of infection so that mortality rate is 49% in subjects entering intensive care units.² It has been reported that critically ill COVID-19 patients remarkably have elevated levels of pro-inflammatory cytokines including IL-6, G-CSF, IP10, MCP1, MIP1A, and TNF α , contributing to outbreak of cytokine storm.³ These turn of events lead to acute respiratory distress syndrome and ultimately maybe death² and glucocorticoids were not efficient in reducing rate of mortality.⁴

It has been suggested that mesenchymal stem cell (MSC) transplantation could be considered as a beneficial approach in treatment many disease. MSC exerts its positive impacts through immunomodulation and differentiation stimulation.⁵ It has proposed that pathogen molecules such as double-stranded RNA in virus is able to increase induction of toll-like receptors (TLRs) on MSC resulting in manifestation of immunomodulatory effects of MSC.⁶ Darwish et al. demonstrated that MSC is potentially able to treat H5N1 infection which has a cytokine profile similar to COVID-19.⁷ The mechanisms

of action of MSC on COVID-19 treatment schematically are shown on Fig. 1.

Therefore, MSC therapy potentially could be considered as an efficacious and safe treatment approach in COVID-19-induced pneumonia. In this regard, Leng et al. recently demonstrated that with perfusion of 1×10^6 cells per kilogram of weight, almost all the clinical symptoms such as fever, breath shortness, and low oxygen saturation disappeared and the inflammation levels were relieved 2–4 days after MSC transplantation.⁸ Moreover, in a critically ill 65 years old woman, MSC transplantation with 5×10^7 cells three times resulted in significant decrease in CRP and increase in CD3+, CD4+ and CD8+ T cells to the normal ranges. Also, CT images implied to remarkable relieve in pneumonia.⁹ Furthermore, there are 30 registered studies investigating MSC therapy on COVID-19 to explore whether MSC transplantation could be able to shed the light in COVID-19 treatment. A summary of characteristics of registered studies are presented in Table 1.

Author contribution

Masoud Khorshidi: Conceptualization, Investigation, Methodology, Software, Visualization; Meysam Zarezadeh: Data curation, Investigation, Validation, Visualization, Writing - original draft, Writing - review and editing; Mohammadreza Emami: Data curation, Visualization; Beheshteh Olang: Data curation, Visualization, Validation; Omid Moradi Moghaddam: Conceptualization, Project administration, Supervision, Visualization.

* Corresponding author at: Trauma and Injury Research Center, Critical Care Medicine Department, Iran University of Medical Sciences, Shahid Hemmat Highway, Tehran, Iran.

** Corresponding author at: Ph.D. of Nutritional Sciences, School of Nutrition and Food Sciences, Tabriz University of Medical Sciences, Attar-Neishaburi St., Golgasht Alley, Azadi Blvd., Tabriz, Iran.

E-mail addresses: zarezhadehm@tbzmed.ac.ir (M. Zarezadeh), moradimoghaddam@yahoo.com (O.M. Moghaddam).